

Dynamic Pupillometry – An indigenous non-invasive screening tool for clinical utility

Siva kumar A V^{a,*}, Padmavathi R^b, Maruthy K N^c, Mahesh Kumar K^d.

^aAssistant Professor, Dept. of Physiology, Narayana Medical College, Nellore, Andhra Pradesh, India.

^bProfessor & Associate Dean – PG Studies, Dept. of Physiology, Sri Ramachandra Institute of Higher Education and Research (SRIHER), Porur, Chennai, Tamilnadu, India

^cProfessor & Head, Dept. of Physiology, Narayana Medical College, Nellore, Andhra Pradesh, India

^dAsst. Professor, Dept. of Physiology & Biochemistry, Govt. Yoga and Naturopathy College & Hospital, Chennai, Tamilnadu, India

ARTICLE INFO

Article history:

Received 10 August 2022

Revised 12 August 2022

Accepted 15 August 2022

Keywords:

Dynamic Pupillometry

Non-invasive screening

Pupillometers

Prognosis

Indigenous

INTRODUCTION

Non-invasive diagnostic practices have been emerging expeditiously in the scientific medical field, which are more preferred, convenient, and effective than invasive procedures. Most of these practices are playing a crucial role in the diagnosis of several clinical diseases using computer-aided imaging devices. The modern non-invasive procedures commonly employed in the clinical sectors are X-ray, Ultrasonography (USG), Computed Tomography (CT), Magnetic Resonance Image (MRI), functional MRI (fMRI) and Positron Emission Tomography (PET)scans [1]. The Pupillometers are one such imaging instrument to measure the pupil size and reactivity. Recently the utility of computerized or portable, handheld pupillometer has been expanding in health care systems [2]. Previous research studies had revealed that pupillometers have neurodiagnostic significance in various clinical and experimental studies [3,4]. The Dynamic Pupillometry (DP) is also gaining empirical significance to evaluate the depiction of diseases and underlying pathophysiological mechanisms involving in the disease by computing Pupillary Light Reflex (PLR). The quantitative pattern of light reflex is a fundamental component of some neurological assessment, and their variables signify the neurological deficits in various diseases and the prognosis in response to an effective treatment regimen [5]. An impaired pupil response may found in neurological deterioration and poor prognosis. Pupillometer is a self-contained camera that objectively measures static pupil diameter and dynamic pupil

response to a standard light stimulus using Infra-Red (IR) videography [6]. The earlier pupillometers in the 1960s were time-consuming, low precision due to the less frame rate. But recently, automated pupillometers are advanced with digital imaging technology, which can provide acceptable pupil measures. However, these pupillometers are too expensive and unaffordable to a regular clinical setup [7].

There is a considerable development in digital imaging tools in the past few years due to the accessibility of cameras with optimal resolution and frame rate. The encroachment of unique optics in these digital cameras conquers hassle-free capturing of good quality images. These cameras can be employed to video graph the pupil reaction under multiple wavelengths of light. The mesopic vision causes pupil dilation captured under Infrared illumination, and photopic vision leads to pupil constriction under bright light illumination [8, 9]. The entire pupil response can be video graphed, which includes maximum dilation to maximum constriction. It can analyze from individual frames of video for pupil diameter using image analysis. Thus a customized simple web camera with an optimum frame rate can record pupil response to light, which provides a quantitative analysis of PLR using infrared videography [10, 11].

Why we need Quantitative measures of PLR?

The pupil is the central aperture of the iris that controls the intensity of the light falling on the retina. The pupils are circular, bilaterally symmetrical, and centric positioned within the iris of healthy individuals. The

*Correspondence at Siva Kumar A V, Assistant Professor, Dept. of Physiology, Narayana Medical College, Nellore, Andhra Pradesh, India..

Email: redhy.sivakumar5@gmail.com

Cite as: Siva Kumar A V et al. Dynamic Pupillometry – An indigenous non-invasive screening tool for clinical utility. The Journal of Medicine and Science 2022; 01(01): 14-16.

© 2022 The Author(s). This is an open access article under the CC-BY. (<https://creativecommons.org/licenses/by/4.0/>)

pupil size varies in adults from 2 – 4 mm in photopic vision and 2 – 8 mm in scotopic vision controlled by two antagonistic muscles, i.e., Sphincter pupillae and dilator pupillae. A bright light stimulus, accommodation reflex reduces pupil size due to the contraction of sphincter pupillae (Miosis) mediated by parasympathetic stimulation. The other stimulants, like cognitive load, attention, alertness, and darkness, cause dilation of the pupil (mydriasis) due to the contraction of dilator pupillae via the sympathetic activation. So the Pupillary light reflex (PLR) examination is a fundamental, comprehensive neurological index in several clinical disorders. The swinging flashlight test (pen torch test) is often employed to evaluate pupil function as a routine clinical examination for bedside patients. However, it does not provide precise quantitative values instead expressed in a qualitative manner (PERLLA – Pupils are round, reactive to light and accommodation), which has a subjective bias, and there is a considerable interobserver variability which leads to inadequate interpretation. The pupil examination of neurosurgical and ICU patients are often essential as they cannot participate in the clinical evaluation due to their severe condition or sedation. In general clinical practice, the subjective pupil examination may be inadequate by stimulus strength, duration of light stimulus surrounding ambience and observer visual acuity, which makes unreliable. So to circumventing the subtle of manual examination of the pupil, the computerized or handheld pupillometers have been employed for objective measurement of pupillary parameters.

Applications of Pupillometry and its role in clinical utility:

The Dynamic Pupillometry indices help in the assessment of autonomic dysfunction [12, 13] as the PLR graphic pattern has two limbs. The downward limb is the constriction phase represents parasympathetic activation to a light stimulus, and the upward limb is the subsequent redilation phase, indicates sympathetic activation after the light exposure. The PLR evaluation is user friendly and convenient for participation than other traditional tests for autonomic assessment. So the lesions in ANS like Addie's tonic pupil, Horner's syndrome, and lesion in the oculomotor nerve can also be found out using PLR. It is also tailored for the diagnosis of cholinergic deficiency syndromes like Parkinson's and Alzheimer's diseases where PLR is altered [14]. The pupillometry also reveals emotions, attention, and evaluation of cognitive disabilities [15,

16]. It is the easiest method than EEG and fMRI for the measuring of prefrontal activity and Visio - spatial discrimination as they are reflected in pupil response and magnitude of PLR [17]. The pupil reactivity also diminished in raise in intracranial pressure after traumatic brain injury, which in turn becomes worst associated with a lack of prognosis [18]. So the PLR has been designated as a prognostic variable in head injury and critical care patients. Pupil monitoring is essential in intensive care unit patients as they reflect the brain stem activity [19]. The abnormal pupil response is also associated with other diagnostic variables like pulse wave velocity [20].

Thus Dynamic pupillometry can be used as a non-invasive screening tool in various clinical evaluations. It is a novel technique that offers a promising approach portable, inexpensive and simple for measurement of the static and dynamic pupil. The Cost effective pupillometer can be made from an ordinary camera which optimum frame rate to videograph and quantify the PLR to interpret the status of clinical condition.

Funding Source

None

Peer-Review

This article has been reviewed by

1. Dr. Moorti A, Associate Professor, Department of Physiology, Coimbatore Medical College and Hospital, Coimbatore, India.
2. Dr. Jeevakala S, Associate Professor, Department of Ophthalmology, Coimbatore Medical College and Hospital, Coimbatore, India.

Declaration of Competing Interest

None

REFERENCES :

1. Kalmár G, Büki A, Kékesi G, Horváth G, Nyúl LG. Image processing-based automatic pupillometry on infrared videos. *Acta Cybernetica*. 2017 Jan 1;23(2):599-613.
2. Larson MD, Behrends M. Portable infrared pupillometry: a review. *Anesthesia & Analgesia*. 2015 Jun 1;120(6):1242-53.
3. Martínez-Ricarte F, Castro A, Poca MA, Sahuquillo J, Expósito L, Arribas M, Aparicio J. Infrared pupillometry. Basic principles and their application in the non-invasive monitoring of neurocritical patients. *Neurología (English Edition)*. 2013 Jan 1;28(1):41-51.

4. Siva kumar AV, Maruthy KN, Maheshkumar K, Padmavathi R. Letter to Editor: Dynamic pupillometer to quantify pupil light reflex (PLR). *European Journal of Ophthalmology*. 2022 May;32(3):NP100-1.
5. Szulewski A, Kelton D, Howes D. Pupillometry as a Tool to Study Expertise in Medicine. *Frontline Learning Research*. 2017;5(3):53-63.
6. Maruthy KN, Padmavathi R, Sowjanya B, MaheshKumar K. Quantitative determination of pupil by dynamic pupillometry using infrared videography–Role in evaluation of autonomic activity. *Clinical Epidemiology and Global Health*. 2020 Sep 1;8(3):728-32.
7. Kumar AS, Padmavathi R, Maruthy KN, Sowjanya B, Kumar K. An Innovative Technique to Evaluate Quantitative Pupillary Light Reflex by Dynamic Pupillometry using Infrared Videography. *Journal of Clinical & Diagnostic Research*. 2019 Apr 1;13(4).
8. Sivakumar AV, Kalburgi-Narayana M, Kuppusamy M, Ramaswamy P, Bachali S. Computerized dynamic pupillometry as a screening tool for evaluation of autonomic activity. *Neurophysiologie Clinique*. 2020 Oct 1;50(5):321-9.
9. Çömez at, kömür b, eser i. Assessment of pupil diameters of emmetropes and myopes under photopic, mesopic and scotopic Conditions, using the infrared pupillometer integrated within Schwind Sirius multifunctional diagnostic device. *Türkiye Klinikleri Tıp Bilimleri Dergisi*. 2012;32(5):1226-34.
10. Kıyloğlu N, Kılıç MA, Kocatürk T, Özkan SB, Bilgen M. A custom-made pupillometer system for characterizing pupillary light response. *Turkish journal of ophthalmology*. 2018 Aug;48(4):185.
11. Nowak W, Żarowska A, Szul-Pietrzak E, Misiuk-Hojło M. System and measurement method for binocular pupillometry to study pupil size variability. *Biomedical engineering online*. 2014 Dec 1;13(1):69.
12. Srinag L, Maruthy KN, Kareem SK, Kumar AS, Kumar K, Gurja JP. Impact of overweight on cognition and psychomotor skills among children with overweight. *Obesity Medicine*. 2020 Jun 1;18:100191.
13. Piha SJ, Halonen JP. Infrared pupillometry in the assessment of autonomic function. *Diabetes research and clinical practice*. 1994 Nov 1;26(1):61-6.
14. Fotiou DF, Stergiou V, Tsiptsios D, Lithari C, Nakou M, Karlovasitou A. Cholinergic deficiency in Alzheimer's and Parkinson's disease: evaluation with pupillometry. *International Journal of Psychophysiology*. 2009 Aug 1;73(2):143-9.
15. Palinko O, Kun AL, Shyrovkov A, Heeman P. Estimating cognitive load using remote eye tracking in a driving simulator. In *Proceedings of the 2010 symposium on eye-tracking research & applications 2010 Mar 22* (pp. 141-144).
16. Kumar AS, Padmavathi R, Mahadevan S, Maruthy KN, Maheshkumar K. Impaired pupillary light reflex indices in Orbital Apex Syndrome—A rare case report. *Journal Français d'Ophthalmologie*. 2021 May 1;44(5):718-22.
17. Ebitz RB, Moore T. Selective modulation of the pupil light reflex by micro stimulation of prefrontal cortex. *Journal of Neuroscience*. 2017 May 10;37(19):5008-18.
18. Jahns FP, Miroz JP, Messerer M, Daniel RT, Taccone FS, Eckert P, Oddo M. Quantitative pupillometry for the monitoring of intracranial hypertension in patients with severe traumatic brain injury. *Critical care*. 2019 Dec 1;23(1):155.
19. MaheshKumar K, Maruthy KN, Padmavathi R. Comparison of photo pulse plethysmography module with Mobil-O-graph for measurement of pulse wave velocity. *Clinical Epidemiology and Global Health*. 2021 Jan 1;9:216-20.
20. Phillips SS, Mueller CM, Nogueira RG, Khalifa YM. A systematic review assessing the current state of automated pupillometry in the NeuroICU. *Neurocritical care*. 2019 Aug 1:1-20.